

NCIN Upper Gastrointestinal Site Specific Clinical Reference Group (UGI SSCRG): Cancer registry analysis

Victoria Coupland, Karen Linklater, Elizabeth Davies

NCIN Upper GI TSSG Clinical Leads Workshop

25th November 2010

Outline

- Incidence of OG and HPB cancers in England between 1998 and 2006.
- Geographic variation in liver, gallbladder and lower oesophageal cancer
- Next steps

Oesophago Gastric cancer groups

• Upper and middle oesophagus	16,216
• Lower oesophagus	31,672
• Oesophagus not otherwise specified	6,633
• Cardia	16,853
• Distal stomach	14,036
• Stomach not otherwise specified	33,586

Hepatic, pancreatic and biliary cancers


• Pancreas	53,265
• Ampulla of Vater	2,855
• Bile duct and biliary tract	10,689
• Liver (excl intrahepatic bile duct)	11,598
• Gallbladder	3,824
• Duodenum	2,248

Geographic variation


- Liver
- Gallbladder
- Lower oesophageal cancer

Liver cancer

Males




Females




Funnel plots: Liver cancer

Males




Females




Funnel plots: Gallbladder cancer

Males





Females



Lower oesophagus



Males

Lower oesophagus




Females

Lower oesophagus



Funnel plots: Lower oesophageal cancer

Males



Females

